

Mathers Clinic

MADRS: Montgomery-Åsberg Depression Rating Scale

Name: _____ Date: _____/_____/_____

Time: _____ : _____ Personal Notes: _____

Place a value for each question in the box. Add all values for the total score.

1. Apparent Sadness

Representing despondency, gloom and despair (more than just ordinary transient low spirits), reflected in speech, facial expression, and posture. Rate by depth and inability to brighten up.

- 0 No sadness.
- 2 Looks dispirited but does brighten up without difficulty.
- 4 Appears sad and unhappy most of the time.
- 6 Looks miserable all the time. Extremely despondent.

2. Reported Sadness

Representing reports of depressed mood, regardless of whether it is reflected in appearance or not.

Includes low spirits, despondency or the feeling of being beyond help and without hope.

- 0 Occasional sadness in keeping with the circumstances.
- 2 Sad or low but brightens up without difficulty.
- 4 Pervasive feelings of sadness or gloominess. The mood is still influenced by external circumstances.
- 6 Continuous or unvarying sadness, misery or despondency.

3. Inner Tension

Representing feelings or ill-defined discomfort, edginess, inner turmoil, mental tension mounting to either panic, dread or anguish. Rate according to intensity, frequency, duration and the extent of reassurance called for.

- 0 Placid. Only fleeting inner tension.
- 2 Occasional feelings of edginess and ill-defined discomfort.
- 4 Continuous feelings of inner tension or intermittent panic which the patient can only master with some difficulty.
- 6 Unrelenting dread or anguish. Overwhelming panic.

4. Reduced Sleep

Representing the experience of reduced duration or depth of sleep compared to the subject's own normal pattern when well.

- 0 Sleeps as usual.
- 2 Slight difficulty dropping off to sleep or slightly reduced, light or fitful sleep.
- 4 Sleep reduced or broken by at least 2 hours.
- 6 Less than 2 or 3 hours sleep.

Continued on the following page.

Adapted from the www.cnsforum.com, MADRS: Montgomery-Åsberg Depression Rating Scale



Mathers Clinic - MADRS: Montgomery-Åsberg Depression Rating Scale

Name: _____ Date: ____/____/____ Time: ____:____:____

5. Reduced Appetite

Representing the feeling of a loss of appetite compared with when well. Rate by loss of desire for food or the need to force oneself to eat.

- 0 Normal or increased appetite.
- 2 Slightly reduced appetite.
- 4 No appetite. Food is tasteless.
- 6 Needs persuasion to eat at all.

6. Concentration Difficulties

Representing difficulties in collecting one's thoughts amounting to an incapacitating lack of concentration. Rate according to intensity, frequency, and degree of incapacity produced.

- 0 No difficulties in concentrating.
- 2 Occasional difficulties in collecting one's thoughts.
- 4 Difficulties in concentrating and sustaining thought which reduces ability to read or hold a conversation.
- 6 Unable to read or converse without great difficulty.

7. Lassitude

Representing difficulty in getting started or slowness in initiating and performing everyday activities.

- 0 Hardly any difficulty in getting started. No sluggishness.
- 2 Difficulties in starting activities.
- 4 Difficulties in starting simple routine activities, which are carried out with effort.
- 6 Complete lassitude. Unable to do anything without help.

8. Inability To Feel

Representing the subjective experience of reduced interest in the surroundings, or activities that normally give pleasure. The ability to react with adequate emotion to circumstances or people is reduced.

- 0 Normal interest in the surroundings and in other people.
- 2 Reduced ability to enjoy usual interests.
- 4 Loss of interest in the surroundings. Loss of feelings for friends and acquaintances.
- 6 The experience of being emotionally paralysed, inability to feel anger, grief or pleasure and a complete or even painful failure to feel for close relatives and friends.

10. Suicidal Thoughts

Representing the feeling that life is not worth living, that a natural death would be welcome, suicidal thoughts, and preparations for suicide. Suicide attempts should not in themselves influence the rating.

- 0 Enjoys life or takes it as it comes.
- 2 Weary of life. Only fleeting suicidal thoughts.
- 4 Probably better off dead. Suicidal thoughts are common, and suicide is considered as a possible solution, but without specific plans or intension.
- 6 Explicit plans for suicide when there is an opportunity. Active preparations for suicide.

Continued on the following page.

Adapted from the www.cnsforum.com, MADRS: Montgomery-Åsberg Depression Rating Scale



Mathers Clinic - MADRS: Montgomery-Åsberg Depression Rating Scale

Name: _____ Date: ____/____/____ Time: _____:

Score For Each Question:

1: _____ 3: _____ 5: _____ 7: _____ 9: _____

2: _____ 4: _____ 6: _____ 8: _____ 10: _____

TOTAL SCORE

Mathers Clinic - MADRS: Montgomery-Åsberg Depression Rating Scale: The Maximum score is 60.

0-6 = Normal

7-19 = Mild Depression

20-34 = Moderate Depression

35-60 = Severe Depression

Adapted from the www.cnsforum.com, MADRS: Montgomery-Åsberg Depression Rating Scale



Crystal Lake: 145 S Virginia St, Crystal Lake, IL 60014

phone: 815.444.9999

fax: **815.986.1363**

Rockford: 6090 Strathmoor Dr, Ste 1, Rockford, IL 61107

phone: 815.444.9999

fax: **815.986.1363**

Woodstock: 715 W Judd St, Woodstock, IL 60098

phone: 815.444.9999

fax: **815.986.1363**

Elgin: 585 N Tollgate Rd, Ste E, Elgin, IL 60123

phone: 847.462.6099

fax: **847.628.6064**

Fox Lake: 81 E Grand Ave, Fox Lake, IL 60020

phone: 224.908.3005

fax: **847.531.1296**